



Contents lists available at ScienceDirect

Marine Pollution Bulletin

journal homepage: www.elsevier.com/locate/marpolbul

Are we eating plastic-ingesting fish?

Daniele de A. Miranda^{a,*}, Gustavo Freire de Carvalho-Souza^{b,c}^a Laboratório de Compostos Orgânicos em Ecossistemas Costeiros e Marinhos (OrganoMAR), Departamento de Oceanografia, Universidade Federal de Pernambuco, Av. Arquitetura, s/n, Cidade Universitária, 50740-550 Recife, Pernambuco, Brazil^b CAPES Foundation, Ministry of Education of Brazil, 70040-020 Brasília, DF, Brazil^c Associação Brasileira do Lixo Marinho (ABLM), Brazil

ARTICLE INFO

Article history:

Received 12 August 2015

Received in revised form 23 November 2015

Accepted 22 December 2015

Available online xxxx

Keywords:

Marine debris

Pellets

Predator fishes

Artisanal fisheries

ABSTRACT

Yes, we are eating plastic-ingesting fish. A baseline assessment of plastic pellet ingestion by two species of important edible fish caught along the eastern coast of Brazil is described. The rate of plastic ingestion by king mackerel (*Scomberomorus cavalla*) was quite high (62.5%), followed by the Brazilian sharpnose shark (*Rhizoprionodon lalandii*, 33%). From 2 to 6 plastic resin pellets were encountered in the stomachs of each fish, with sizes of from 1 to 5 mm, and with colors ranging from clear to white and yellowish. Ecological and health-related implications are discussed and the potential for transferring these materials through the food-chain are addressed. Further research will be needed of other species harvested for human consumption.

© 2015 Published by Elsevier Ltd.

Marine debris is a growing threat to marine wildlife (Moore et al., 2001). Plastic pollution is found in all coastal and marine environments, from beaches, reefs, mangroves and estuaries, to the open ocean, and takes on many different shapes and forms (pellets, debris, and objects) (Ivar do Sul et al., 2009; Carvalho-Souza and Tinôco, 2011; Costa et al., 2009; Mordecai et al., 2011; Vieira et al., 2011).

Several papers have explored issues related to the accumulation and distribution of plastics in marine habitats and the resulting impacts on aquatic species and various ecological processes (e.g., behavior, bio-invasion, food resources, chemical pollutants) (Laist, 1997; Barnes et al., 2009; Gregory, 2009; Browne et al., 2011; Rochman et al., 2014; Carvalho-Souza, 2015). The ingestion of marine debris by marine wildlife, ranging from zooplankton to marine megafauna (fish, seabirds, sea turtles, and marine mammals) has been widely documented (Laist, 1997; Tourinho et al., 2010; Schuyler et al., 2013; Di Benedetto and Awabdi, 2014).

The first reports of marine debris ingestion by fish was published by Carpenter et al. (1972) and described the presence of plastic particles in larvae and adult fish. Many other species of fish, rays, and sharks have been documented ingesting plastic debris in recent decades, with the number of records still growing steadily (Hoss and Settle, 1990; Laist, 1997; Jackson et al., 2000; Cliff et al., 2002; Boerger et al., 2010; Possatto et al., 2011; Dantas et al., 2012; Choy and Drazen, 2013; Jantz et al., 2013; Foekema et al., 2013; Lusher et al., 2013; Di Benedetto and Awabdi, 2014; Ramos et al., 2012; Romeo et al., 2015).

Plastic can cause direct damage to marine wildlife through entanglement, malnutrition (gut blockage and pseudo-satiation), suffocation and decreased mobility – often resulting in their death (Laist, 1997). Indirect effects have also been documented relating to the accumulation of heavy metals and chemical pollutants such as polychlorinated biphenyls (PCBs), polybrominated diphenyls (PBDEs), and polycyclic aromatic hydrocarbons (PAHs) (Teuten et al., 2009; Frias et al., 2010).

Other harmful effects of discarded plastics include the transport of alien/invasive species (so-called 'hitch-hiking') and the inhibition of gas-exchange and the resulting smothering of seabeds (Gregory, 2009).

Rochman and Browne (2013) proposed classifying plastics (synthetic polymers) as hazardous wastes, and the risks of plastic ingestion are now focusing on its transfer between trophic levels, health impacts across food-chains, and possible effects on humans. New research is needed in this key area.

Laboratory studies have confirmed plastic transfer between mussels and crabs (Farrel and Nelson, 2013); a more recent investigation reported the ingestion of plastic microspheres by several zooplankton groups and plastic micro-particle transfer (via planktonic organisms) between trophic levels (mesozooplankton-macrozooplankton) (Setälä et al., 2014). Likewise, medaka fish exposed to a complex mixture of plastic (and their associated chemical pollutants) via ingestion showed disturbances of endocrine system functioning (Rochman et al., 2013, 2014). Matsson et al. (2015) – demonstrating that nanoparticle uptake occurs throughout the algae-zooplankton-fish food chain with resulting genetic, morphological, and behavioral alterations.

In light of these disturbing impacts, it will be important to identify fish species showing plastic residues in their stomachs. We report here a baseline assessment of the ingestion of plastic pellets by two

* Corresponding author at: UFPE, 50740-550 Recife, PE, Brazil.

E-mail addresses: danielemirandaba@hotmail.com (D.A. Miranda), gustavofscouza@yahoo.com.br (G.F. de Carvalho-Souza).

commercially valuable fish species caught by artisanal fishermen on the eastern coast of Brazil.

The study area was located along the coast of the city of Salvador, Bahia State, in northeastern Brazil (Fig. 1).

Salvador is the capital of Bahia State and the third largest city in Brazil, with an estimated population of nearly 3 million (IBGE, 2014). The city has two public ports, seven private marine terminals, one oil refinery, and the largest integrated industrial-petrochemical complex in Latin America (since the late 1970s).

The plastic production chains in these industries are composed of three basic steps: 1) monomer generation (cracking, e.g., ethene, propene, butene, benzene); 2) polymerization (e.g., the formation of polyethylene, polypropylene, polystyrene, PVC, EVA, and others); and 3) transformation (manufacturing finished products for consumer markets) (ABIPLAST, 2013).

Occasional unintentionally resin pellet losses into rivers and other watercourses can occur during these steps or during logistic stages (e.g., terrestrial/aquatic transport), and these plastics will eventually migrate to coastal and marine environments (Ogata et al., 2009).

Metal and solid waste contamination has been documented in the Todos os Santos Bay in the metropolitan region of Salvador, impacting beaches, rocky shores, estuaries, and mangrove swamps (Hatje et al., 2010; Carvalho-Souza and Tinoco, 2011; Eça et al., 2013). These sites are known to be nurseries for many aquatic species and are used by artisanal fishermen. Coastal fisheries are an important economic activity in many areas in northeastern Brazil (Diegues, 2008).

Sampling was undertaken from April to May/2011 at two fishing ports in the city of Salvador: Pituba (Z1-APEPI) and Itapuã (Z6-COOPI) (Fig. 1). These are traditional fishing communities (e.g., COOPI was established more than two centuries ago) with artisanal fleets of small wooden boats. The fishermen generally use hook and line

fishing techniques to capture high-value specimens (Dominguez et al., 2013).

Fish were collected randomly at the landing points from among recently captured specimens. Biometric measurements (furcal length – cm, weight – g) and taxonomic identifications (based on the scientific literature – Carvalho-Filho, 1999; Froese and Pauly, 2014) were performed at the landing points. The stomachs were removed from the fish and subsequently examined in the laboratory using a binocular dissecting microscope (40×).

The stomach contents were analyzed and separated into natural foods (prey and organic material) and plastic items. The plastic debris was further separated, quantified, and then classified according to its shapes and colors. The fish species were sorted according to their habitats (pelagic or demersal) and trophic categories (Carvalho-Filho, 1999; Ferreira et al., 2004; Bornatowski et al., 2012; Froese and Pauly, 2014), noting, for each individual, where it was captured, the presence of plastics debris, and the numbers of plastic items and their frequencies.

The stomach contents of 32 fish specimens belonging 11 species and 9 families (2 elasmobranches and 9 teleosts) were examined (Table 1). Resin pellets were the only type of plastic observed in their stomachs. Plastic pellets were found in the stomachs of 7 individuals (22%) of two species: the king mackerel, *Scomberomorus cavalla* (Cuvier, 1829) (Scombridae) and the Brazilian sharpnose shark, *Rhizoprionodon lalandii* (Müller & Henle, 1839) (Carcharhinidae) (Fig. 2a–b). The highest frequency of plastic occurrence was recorded for *S. cavalla*, with 5 specimens containing pellets (62.5% of the individuals of the species), followed by 2 individuals of *R. lalandii* (33%). From 2 to 6 plastic pellets were encountered in *S. cavalla*, with sizes ranging from 2 to 5 mm in their longest dimension. From 1 to 3 small pellets were encountered in individuals of *R. lalandii* with sizes varying from 1 to 3




Fig. 1. Map of the metropolitan region of Salvador indicating two fishing ports, Bahia State, Brazil.

Table 1

The fish species analyzed with quantitative data concerning the micro-plastic pellets ingested.

Taxa	N Fish	N Fish Plastic	FOP (%)	N Plastic	Size Pellet (mm)	Habitat and Trophic Categories
<i>Acanthurus coeruleus</i> (Bloch & Schneider, 1801)	1	-	-	-	-	Reef/Roving Herbivore
<i>Caranx crysos</i> (Mitchill, 1815)	3	-	-	-	-	Pelagic-Coastal/Carnivore
<i>Dasyatis americana</i> (Hildebrand & Schroeder, 1928)	3	-	-	-	-	Sandy and Muddy Bottoms/Carnivore
<i>Lagocephalus laevigatus</i> (Linnaeus, 1766)	1	-	-	-	-	Pelagic-Coastal, Sandy and Muddy Bottoms/Omnivore
<i>Lutjanus analis</i> (Cuvier, 1828)	2	-	-	-	-	Reef/Carnivore
<i>Lutjanus jocu</i> (Cuvier, 1828)	5	-	-	-	-	Reef/Carnivore
<i>Mycteroperca</i> sp.	1	-	-	-	-	Reef/Carnivore
<i>Paralichthys brasiliensis</i> (Agassiz, 1831)	1	-	-	-	-	Sandy and Muddy Bottoms/Omnivore
<i>Rhizoprionodon landii</i> (Müller & Henle, 1839)	6	2	62.5	2-Jun	1-Mar	Pelagic-demersal/Carnivore
<i>Scomberomorus cavalla</i> (Cuvier, 1829)	8	5	33	1-Mar	2-May	Benthic-pelagic/Carnivore
<i>Sphyrna guachancho</i> (Cuvier, 1829)	1	-	-	-	-	Pelagic-Coastal/Carnivore

mm. The plastic resin pellets encountered had cylindrical shapes, and colors ranging from clear to white, and yellowish.

This study quantified the incidence of plastic pellets in the stomachs of carnivorous fish considered “generalist-predators” species that occupy high trophic levels. These predator species play a key role in environmental dynamics and contribute to maintaining the overall balance of marine ecosystems (Heithaus et al., 2008). Intensive fishing, habitat losses, and pollution have caused serious declines in the populations of large marine fish, and the present records are apparently the first to describe plastic pellet ingestion in two commercially important fish species (*S. cavalla* and *R. landii*) along the Brazilian coast.

The king mackerel is one of the most important fishery resources caught by artisanal fishermen in northeastern Brazil (Lessa et al., 2004). *S. cavalla* occurs in the Western Atlantic from Massachusetts (USA) south to Sao Paulo State (Brazil) (ICCAT, 2014). This species is considered over-exploited in the Gulf of Mexico, and protective actions are recommended for northeastern Brazil to avoid over-fishing (Lessa et al., 2004).

R. landii is an abundant, small, coastal elasmobranch distributed from the southwestern to western-central Atlantic Ocean (Rosa et al., 2004). The Brazilian sharpnose shark is important commercial resource commonly caught by artisanal fishermen; it has been reported to be impacted by marine pollution in the form of entanglement in plastic collars (Sazima et al., 2002; Rosa et al., 2004).

Fish have been consumed since time immemorial by human societies, and are known for their many health benefits (McManus and Newton, 2011). Organic compounds (e.g., PAHs) and heavy metals are known to impact fish behavior and health (Atchison et al., 1987; Ajiboye et al., 2011; Sevcikova et al., 2011; Weis and Candelmo, 2012), and maximum contamination levels have been established by government health agencies worldwide (Commission Regulation, 2006; Buchman, 2008; Brasil, 2013).

Many concerns regarding toxic compound transfers between trophic levels have resulted in laboratory experiments showing severe impacts of plastics on marine wildlife – including endocrine disruption and behavioral, physiological, and metabolic alterations (Rochman et al., 2013, 2014; Matsson et al., 2015).

Plastic pellets are a clear threat in aquatic environments due to their potential for accumulating and transporting contaminants adsorbed from the surrounding water (Teuten et al., 2009; Ogata et al., 2009). Carpenter et al. (1972) found white polyethylene spherules in the guts of several fish species caught along the New England coast (USA), indicating probable consumption selectivity. The three pellet colors (clear, white, and yellowish) encountered in the present work suggest that these plastics had been adrift in the sea for long periods of time and had undergone oxidation (Ogata et al., 2009) – although we cannot discount the possibility that digestive enzymes in the fishes' stomachs provoked color alterations. According to Endo et al. (2005), yellowing pellets may contain high concentrations of persistent organic pollutants (POPs).

Aquatic contamination by plastic pellets has been well documented globally (Ogata et al., 2009) and Brazil is not exempt (Costa et al., 2009; Ivar do Sul et al., 2009; Fisner et al., 2013; Turra et al., 2014). According to the International Pellet Watch (IPW), high concentrations of DDT have been identified in pellet samples collected in the waters off southern Bahia State (IPW, 2014).

The percentage of predator fish with ingested plastic pellets (50 %) was similar to, or greater than, reports of other studies of pelagic and demersal species. Lusher et al. (2013), studying pelagic and demersal fish in the English Channel, found plastic residues in 36.5% of the gastrointestinal tracts examined. The stomach contents of the sharpnose lancetfish, *Alepisaurus ferox*, were examined in the northern Pacific, and 24.5% of all individuals were found to contain plastic marine debris (Jantz et al., 2013). Pelagic predatory fish were reported to have frequencies of plastic ingestion ranging from <1% to 58% in the central-northern Pacific. Plastic ingestion has been documented in estuarine fishes such as catfish (23%), drums (7.9%), and gerreids (13.4%) (Possatto et al., 2011; Dantas et al., 2012; Ramos et al., 2012) as well as in pelagic coastal fish (0.7%) in Brazil (Di Benedetto and Awabdi, 2014). Even more recently, Romeo et al. (2015) encountered macro-, meso-, and micro-plastic debris in 12.5% to 32.4% of the stomachs of three large pelagic fish (*Xiphias gladius*, *Thunnus thynnus*, and *Thunnus alalunga*) in the Mediterranean Sea.




Fig. 2. Samples of the plastic pellets encountered in the stomach contents of fish. Scale (cm).

Plastic ingestion by marine organisms is commonly reported as “mistaken identity” consumption because of its similarity to natural prey. This hypothesis is well supported in the case of sea turtles ingesting clear plastic items (e.g., plastic bags floating in the water) due to their 3D shapes and movements that are similar to jellyfish (Schuyler et al., 2013). Many species of seabirds have also been found with high concentrations of micro-plastic (transparent or colored) debris in their digestive tracts, which were apparently mistaken for potential prey (e.g., fish eggs, plankton) (Azzarello and Van-Vleet, 1987; Laist, 1997). Nylon ingestion by estuarine fish during suction feeding in sediments could reflect their resemblance to polychaetes (Possatto et al., 2011) or to the consumption of organisms aggregated to plastic debris (mixed with natural prey items) (Ramos et al., 2012). These hypotheses do not appear appropriate for explaining the ingestion of micro-plastic debris by top fish predators.

Recent hypotheses concerning the consumption of plastic debris by large fish have considered different aspects of their predatory feeding behaviors, including opportunist strategies, accidental ingestion during feeding strikes, or the ingestion of aggregated prey items (Battaglia et al., 2013; Romeo et al., 2015).

Previous studies of plastic ingestion, and available information on the dietary and feeding ecology of *S. cavalla* and *R. lalandii*, provide clues and suggest two hypotheses concerning why these “generalist-predator” species also ingest micro-plastic debris (Fig. 3).

- (i) The voracious feeding habits of generalist predatory fish result in the non-intentional ingestion of plastic debris. Mackerel fish (genus *Scomberomorus*) are known to perform rapid feeding attacks in pelagic environments, during vertical migrations, and on the seabed (Godcharles and Murphy, 1986; ICCAT, 2014). Similarly, sharpnose sharks (genus *Rhizoprionodon*) are opportunistic coastal predators dwelling and foraging along sandy

and muddy inshore bottoms (Silva and Almeida, 2001; Rosa et al., 2004; Bornatowski et al., 2012). Some predator fishes (e.g., sharks) are also known to tentatively bite objects to examine them, and have been observed attacking drifting plastic (see Carson, 2013). As plastic micro-pellets are commonly encountered floating in aquatic environments or mixed with surface sediments (Ivar do Sul and Costa, 2013), there ingestion can occur non-intentionally during foraging and attack behavior.

- (ii) Ingestion can occur through the food chain if micro-plastics were ingested by smaller prey items.

Fish such as clupeids and scianids (both reported ingesting plastic, see Carpenter et al., 1972; Dantas et al., 2012), together with crustaceans and squids, are the main food items of larger generalist-predators, (Bornatowski et al., 2012; ICCAT, 2014). Cephalopods and crustaceans are known to ingest plastic debris (Murray and Cowie, 2011; Braid et al., 2012), and the food-chain hypothesis has been described for marine mammals (Eriksson and Burton, 2003).

The ingestion of plastic pellets by predatory fish therefore comes as no surprise, but indicates that this type of contamination in tropical coastal areas will certainly increase and will require more attention in the near future – especially in terms of human health – as resin pellets in the guts of fish prized for human consumption are a potential public health hazard.

Environmental pollution by plastics is a growing challenge for health and governmental agencies, society as a whole, and the scientific community. Further studies will be needed of commercially valuable species destined for human consumption, to verify their health statuses, maximum contamination tolerance levels, and the potential transfer of contaminants to upper trophic levels in the environment – including humans.




Fig. 3. Conceptual model of plastic pellet ingestion by generalist-predators.

Acknowledgments

The authors wish to thank UCSal for logistic support, the artisanal fishing communities (Pituba and Itapuã), and especially “Boi” and his brothers, and “Pai Velho” and collaborators. We are grateful to R. Funch for reviewing the English text. D. Miranda acknowledges and thanks CNPq, and GF Carvalho-Souza thanks CAPES for their scholarships.

References

- ABIPLAST, 2013. Perfil 2013. Brazilian Plastic Processed Industry, Sao Paulo, Brazil, p. 33.
- Ajiboye, O.O., Yakubu, A.F., Adams, T.E., 2011. A review of polycyclic aromatic hydrocarbons and heavy metal contamination of fish from fish farms. *J. Appl. Sci. Environ. Manag.* 15 (1), 235–238.
- Atchison, G.J., Henry, M.G., Sandheinrich, M.B., 1987. Effects of metals on fish behavior: A review. *Environ. Biol. Fish* 18, 11–25.
- Azzarello, M.Y., Van-Vleet, E.S., 1987. Marine birds and plastic pollution. *Mar. Ecol. Prog. Ser.* 37, 295–303.
- Barnes, D.K.A., Galgani, F., Thompson, R.C., Barlaz, M., 2009. Accumulation and fragmentation of plastic debris in global environments. *Philos. Trans. R. Soc. B* 364, 1985–1998.
- Battaglia, P., Andaloro, F., Consoli, P., Esposito, V., Malara, D., Musolino, S., Pedà, C., Romeo, T., 2013. Feeding habits of the Atlantic bluefin tuna, *Thunnus thynnus* (L. 1758), in the central Mediterranean Sea (Strait of Messina). *Helgol. Mar. Res.* 67 (1), 97–107.
- Boerger, C.M., Lattin, G.L., Moore, S.L., Moore, C.J., 2010. Plastic ingestion by planktivorous fishes in the North Pacific Central Gyre. *Mar. Pollut. Bull.* 60, 2275–2278.
- Bornatowski, H., Heithaus, M.R., Abilhoa, V., Corrêa, M.F.M., 2012. Feeding of the Brazilian sharpnose shark *Rhizoprionodon lalandii* (Müller & Henle, 1839) from southern Brazil. *J. Appl. Ichthyol.* 28, 623–627.
- Braid, H.E., Deeds, J., DeGrasse, J.L., Wilson, J.J., Osborne, J., Hanner, R.H., 2012. Preying on commercial fisheries and accumulating paralytic shellfish toxins: a dietary analysis of invasive *Dosidicus gigas* (Cephalopoda Ommastrephidae) stranded in Pacific Canada. *Mar. Biol.* 159, 25–31.
- BRASIL, 2013. Ministério da Saúde. Resolução n°42, de 30 de agosto de 2013. Agência Nacional de Vigilância Sanitária (<<http://portal.anvisa.gov.br/>> accessed 20.01.14).
- Browne, M.A., Crump, P., Niven, S.J., Teuten, E., Tonkin, A., Galloway, T., Thompson, R., 2011. Accumulation of microplastic on shorelines worldwide: sources and sinks. *Environ. Sci. Technol.* 45, 21.
- Buchman, M.F., 2008. National Oceanic and Atmospheric Administration (NOAA). Screening Quick Reference Tables. (SQIRTS) (<<http://archive.orr.noaa.gov/>> accessed 01.20.14).
- Carpenter, E.J., Anderson, S.J., Harvey, G.R., Miklas, H.P., Peck, B.B., 1972. Polystyrene spherules in coastal waters. *Science* 178, 749–750.
- Carson, H.S., 2013. The incidence of plastic ingestion by fishes: from the prey's perspective. *Mar. Pollut. Bull.* 74, 170–174.
- Carvalho-Filho, A., 1999. Peixes da costa brasileira. 3. ed. Melro, São Paulo (318 pp.).
- Carvalho-Souza, G.F., 2015. Association between juvenile carangid fish with the jellyfish *Aurelia* sp. and a floating plastic bag. *J. Mar. Biol. Oceanogr.* 4, 1.
- Carvalho-Souza, G.F., Tinoco, M.S., 2011. Avaliação do lixo marinho em costões rochosos na Baía de Todos os Santos, Bahia, Brasil. *Rev. Gestão Costeira Integrada* 11 (1), 135–143.
- Choy, C.A., Drazen, J.C., 2013. Plastic for dinner? Observations of frequent debris ingestion by pelagic fishes from the central North Pacific. *Mar. Ecol. Prog. Ser.* 485, 155–163.
- Cliff, G., Dudley, S.F.J., Ryan, P.G., Singleton, N., 2002. Large sharks and plastic debris in KwaZulu-Natal, South Africa. *Mar. Freshw. Res.* 53, 575–581.
- Commission Regulation, 2006. (EC) No 1881/2006 of 19 December 2006 setting maximum levels for certain contaminants in foodstuffs (OJ L 364, 20.12.2006, p. 5). (<<http://eur-lex.europa.eu/legal-content/ES/AL/?uri=CELEX:32006R1881>> accessed 01.20.14.).
- Costa, M.F., Ivar do Sul, J.A., Silva-Cavalcanti, J.S., Araújo, M.C.B., Spengler, A., Tourinho, P.S., 2009. On the importance of size of plastic fragments and pellets on the strandline: a snapshot of a Brazilian beach. *Environ. Monit. Assess.* 168 (1–4), 299–304.
- Dantas, D.V., Barletta, M., Costa, M.F., 2012. The seasonal and spatial patterns of ingestion of polyfilament nylon fragments by estuarine drums (Sciaenidae). *Environ. Sci. Pollut. Res.* 19, 600–606.
- Di Benedetto, A.P.M., Awabdi, D.R., 2014. How marine debris ingestion differs among megafauna species in a tropical coastal area. *Mar. Pollut. Bull.* 88, 86–90.
- Diegues, A., 2008. Marine protected areas and artisanal fisheries in Brazil. *Samudra monographs. International Collective in Support of Fishworkers, Chennai*.
- Dominguez, J.M.L., Silva, R.P., Nunes, A.S., Freire, A.F.M., 2013. The narrow, shallow, low-accommodation shelf of central Brazil: Sedimentology, evolution, and human uses. *Geomorphology* 203, 46–59.
- Eça, G.F., Pedreira, R.M.A., Hatje, V., 2013. Trace and major elements distribution and transfer within a benthic system: Polychaete *Chaetopterus variopedatus*, commensal crab *Polonyx gibbesi*, worm tube, and sediments. *Mar. Pollut. Bull.* 74, 32–41.
- Endo, S., Takizawa, R., Okuda, K., Takada, H., Chiba, K., Kanehiro, H., Ogi, H., Yamashita, R., Date, T., 2005. Concentration of polychlorinated biphenyls (PCBs) in beached resin pellets: variability among individual particles and regional differences. *Mar. Pollut. Bull.* 50, 1103–1114.
- Eriksson, C., Burton, H., 2003. Origins and biological accumulation of small plastic particles in fur seals from Macquarie Island. *Ambio* 32, 380–384.
- Farrel, P., Nelson, K., 2013. Trophic level transfer of microplastic: *Mytilus edulis* (L.) to *Carcinus maenas* (L.). *Environ. Pollut.* 177, 1–3.
- Ferreira, C.E.L., Floeter, S.R., Gasparine, J.L., Joyeux, J.C., Ferreira, B.P., 2004. Trophic structure patterns of Brazilian reef fishes: a latitudinal comparison. *J. Biogeogr.* 31, 1093–1106.
- Fisner, M., Taniguchi, S., Moreira, F., Bicego, M.C., Turra, A., 2013. Polycyclic aromatic hydrocarbons (PAHs) in plastic pellets: Variability in the concentration and composition at different sediment depths in a sandy beach. *Mar. Pollut. Bull.* 70, 219–226.
- Foekema, E.M., De Groot, C., Mergia, M.T., Murk, A.J., van Franeker, J.A., Koelmans, A.A., 2013. Plastic in North Sea fish. *Environ. Sci. Technol.* 47, 8818–8824.
- Frias, J.P.G.L., Sobral, P., Ferreira, A.M., 2010. Organic pollutants in microplastics from two beaches of the Portuguese coast. *Mar. Pollut. Bull.* 60, 1988–1992.
- Froese, R., Pauly, D., 2014. FishBase. World Wide Web electronic publication (Available at <<http://www.fishbase.org>> accessed 20.01.14).
- Godcharles, M.F., Murphy, M.D., 1986. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (south Florida) - king mackerel and Spanish mackerel. U. S. Fish Wildl. Serv. Biol. Rep. 82(11.58). U. S. Army Corps of Engineers (TR EL-82-4. 18).
- Gregory, M.R., 2009. Environmental implications of plastic debris in marine settings entanglement, ingestion, smothering, hangers-on, hitch-hiking and alien invasions. *Philos. Trans. R. Soc. B* 364, 2013–2025.
- Hatje, V., Macedo, S.M., de Jesus, R.M., Cotrim, G., Garcia, K.S., de Queiroz, A.F., Ferreira, S.L.C., 2010. Inorganic as speciation and bioavailability in estuarine sediments of Todos os Santos Bay, BA, Brazil. *Mar. Pollut. Bull.* 60, 2225–2232.
- Heithaus, M.R., Frid, A., Wirsing, A.J., Worm, B., 2008. Predicting ecological consequences of marine top predator declines. *Trends Ecol. Evol.* 23, 202–210.
- Hoss, D.E., Settle, L.R., 1990. In: Shomura, R.S., Godfrey, M.L. (Eds.), Ingestion of plastic by teleost fishes. Proceedings of the Second International Conference on Marine Debris 2–7 April 1989, Honolulu, Hawaii, U.S. Department of Commerce, NOAA Tech. Memo. NMFS, NOAA-TM-NMFS-SWFC-154, pp. 693–709.
- IBGE, 2014. Diretoria de Pesquisas, Coordenação de População e Indicadores Sociais. (Available at <<http://www.cidades.ibge.gov.br>> accessed 20.01.14).
- ICCAT, 2014. ICCAT Manual. International Commission for the Conservation of Atlantic Tuna. ICCAT Publications [on-line] Updated 2014. [Cited 01/27/]. <http://www.iccat.int/en/ICCATManual.htm>.
- IPW, 2014. International Pellet Watch. Global Monitoring of Persistent Organic Pollutants (POPs) using Beached Plastic Resin Pellets. DDTs: Global Pollution Map, Brazil (Available at, <<http://www.pelletwatch.org/maps/map-2.html#Brazil>>, accessed 20.01.14).
- Ivar do Sul, J.A., Costa, M.F., 2013. The present and future of microplastic pollution in the marine environment. *Environ. Pollut.* S0269-7491 (13).
- Ivar do Sul, J.A., Spengler, A., Costa, M.F., 2009. Here, there and everywhere. Small plastic fragments and pellets on beaches of Fernando de Noronha (Equatorial Western Atlantic). *Mar. Pollut. Bull.* 58, 1229–1244.
- Jackson, G.D., Buxton, N.G., George, M.J.A., 2000. Diet of the southern opah *Lampris immaculatus* on the Patagonian Shelf: the significance of the squid *Moroteuthis ingens* and anthropogenic plastic. *Mar. Ecol. Prog. Ser.* 206, 261–271.
- Jantz, L.A., Morishige, C.L., Bruland, G.L., Lepczyk, C.A., 2013. Ingestion of plastic marine debris by sharpnose lancefish (*Alepisaurus ferox*) in the North Pacific Ocean. *Mar. Pollut. Bull.* 69, 97–104.
- Laist, D.W., 1997. Impacts of marine debris: entanglement of marine life in marine debris including a comprehensive list of species with entanglement and ingestion records. In: Coe, J.M., Rogers, D.B. (Eds.), Marine Debris: sources, impacts, and solutions. Springer Series on Environmental Management, New York.
- Lessa, R.P., Nóbrega, M.F., Bezerra Jr., J.L., 2004. Dinâmica de populações e avaliação de estoques de recursos pesqueiros na Região Nordeste. Programa de avaliação do potencial sustentável de recursos vivos na zona econômica Exclusiva (REVIZEE) Relatório Síntese 274. Subcomitê Regional Nordeste (Score-NE), Recife, PE.
- Lusher, A.L., McHugh, M., Thompson, R.C., 2013. Occurrence of microplastics in the gastrointestinal tract of pelagic and demersal fish from the English Channel. *Mar. Pollut. Bull.* 67, 94–99.
- Matsson, K., Eklvall, M.T., Hansson, L.A., Linse, S., Anders, M., Cedervall, T., 2015. Altered Behavior, Physiology, and Metabolism in Fish Exposed to Polystyrene Nanoparticles. 49 (1), 553–561 (†).
- McManus, A., Newton, W., 2011. Seafood, nutrition and human health: A synopsis of the nutritional benefits of consuming seafood. Centre of Excellence Science, Seafood & Health, Curtin Health Innovation Research Institute, Curtin University of Technology, Perth.
- Moore, C.J., Moore, S.L., Leecaster, M.K., Weisberg, S.B., 2001. A comparison of plastic and plankton in the North Pacific Central Gyre. *Mar. Pollut. Bull.* 42, 1297–1300.
- Mordecai, G., Tyler, P.A., Masson, D.G., Huvenne, V.A.L., 2011. Litter in submarine canyons off the west coast of Portugal. *Deep Sea Res. II Top. Stud. Oceanogr.* 58 (23–24), 2489–2496.
- Murray, F., Cowie, P.R., 2011. Plastic contamination in the decapod crustacean *Nephrops norvegicus* (Linnaeus, 1758). *Mar. Pollut. Bull.* 62, 1207–1217.
- Okata, Y., Takada, H., Mizukawa, K., Hiraia, H., Iwasawa, S., Endo, S., Mato, Y., Saha, M., Okuda, K., Nakashima, A., Murakami, M., Zurcher, N., Booyatumanondo, R., Zakaria, M.P., Dung, L.Q., Gordon, M., Miguez, C., Suzuki, S., Moore, C., Karapanagiotik, H.K., Weerts, S., McClurg, T., Burres, E., Smith, W., Van Velkenburg, M., Lang, J.S., Lang, R.C., Laursen, D., Danner, B., Stewardson, N., Thompson, R.C., 2009. International Pellet Watch: global monitoring of persistent organic pollutants (POPs) in coastal waters. 1. Initial phase data on PCBs, DDTs, and HCHs. *Mar. Pollut. Bull.* 58, 1437–1446.
- Possatto, F.E., Barletta, M., Costa, M.F., do Sul, J.A., Dantas, D.V., 2011. Plastic debris ingestion by marine catfish: an unexpected fisheries impact. *Mar. Pollut. Bull.* 62, 1098–1102.
- Ramos, J.A.A., Barletta, M., Costa, M.F., 2012. Ingestion of nylon threads by Gerreidae while using a tropical estuary as foraging grounds. *Aquat. Biol.* 17, 29–34.
- Rochman, C.M., Browne, M.A., 2013. Classify plastic waste as hazardous. *Nature* 494 (7436), 169–171.

- Rochman, C.M., Hoh, E., Kurobe, T., Teh, S.J., 2013. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. *Sci. Rep.* 3, 3263.
- Rochman, C.M., Kurobe, T., Teh, S.J., 2014. Early warning signs of endocrine disruption in adult fish from the ingestion of polyethylene with and without sorbed chemical pollutants from the marine environment. *Sci. Total Environ.* 493, 656–661.
- Romeo, T., Pietro, B., Pedà, C., Consoli, P., Andaloro, F., Fossi, M.C., 2015. First evidence of presence of plastic debris in stomach of large pelagic fish in the Mediterranean Sea. *Mar. Pollut. Bull.* 95, 358–361.
- Rosa, R.S., Gadig, O.B.F., Santos Motta, F., Namora, R.C., 2004. *Rhizoprionodon lalandii*. The IUCN Red List of Threatened Species (Version 2014.3 <www.iucnredlist.org>. Downloaded on 03 February 2015).
- Sazima, I., Gadig, O.B.F., Namora, R.C., Motta, F.S., 2002. Plastic debris collars on juvenile carcharhinid sharks (*Rhizoprionodon lalandii*) in southwest Atlantic. *Mar. Pollut. Bull.* 44, 1147–1149.
- Schuyler, Q., Hardesty, B.D., Wilcox, C., Townsend, K., 2013. Global analysis of anthropogenic debris ingested by sea turtles. *Conserv. Biol.* 28, 129–139.
- Setälä, O., Fleming-Lehtinen, V., Lehtiniemi, M., 2014. Ingestion and transfer of microplastics in the planktonic food web. *Environ. Pollut.* 185, 77–83.
- Sevcikova, M., Modra, H., Slaninova, A., Svobodova, Z., 2011. Metals as a cause of oxidative stress in fish: a review. *Vet. Med.* 11, 537–546.
- Silva, C.M.L., Almeida, Z.S., 2001. Alimentação de *Rhizoprionodon porosus* (Elasmobranchii: Carcharhinidae) da costa do Maranhão, Brasil. *Bol. Inst. Pesca* 27, 201–207.
- Teuten, E.L., Saquing, J.M., Knappe, D.R., Barlaz, M.A., Jonsson, S., Bjorn, A., Rowland, S.J., Thompson, R.C., Galloway, T.S., Yamashita, R., Ochi, D., Watanuki, Y., Moore, C., Viet, P.H., Tana, T.S., Prudente, M., Boonyatumanond, R., Zakaria, M.P., Akkhavong, K., Ogata, Y., Hirai, H., Iwasa, S., Mizukawa, K., Hagino, Y., Imamura, A., Saha, M., Takada, H., 2009. Transport and release of chemicals from plastics to the environment and to wildlife. *Philos. Trans. R. Soc. Lond. Ser. B Biol. Sci.* 364, 2027–2045.
- Tourinho, P.S., Fillmann, G., Ivar do Sul, J.A., 2010. Is marine debris ingestion still a problem for the coastal marine biota of southern Brazil? *Mar. Pollut. Bull.* 60, 396–401.
- Turra, A., Manzano, A.B., Dias, R.J.S., Mahiques, M.M., Barbosa, L., Balthazar-Silva, D., Moreira, F.T., 2014. Three-dimensional distribution of plastic pellets in sandy beaches: shifting paradigms. *Sci. Rep.* 4. <http://dx.doi.org/10.1038/srep04435>.
- Vieira, B.P., Dias, D., Hanazaki, N., 2011. Homogeneidade de Encalhe de Resíduos Sólidos em um Manguezal da Ilha de Santa Catarina, Brasil. *Rev. Gestão Costeira Integr.* 11 (1), 21–30.
- Weis, J.S., Candelmo, A., 2012. Pollutants and fish predator/prey behavior: a review of laboratory and field approaches. *Curr. Zool.* 58, 9–20.